



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2494179 A1 2004/02/19

(21) 2 494 179

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2003/08/07
(87) Date publication PCT/PCT Publication Date: 2004/02/19
(85) Entrée phase nationale/National Entry: 2005/02/04
(86) N° demande PCT/PCT Application No.: JP 2003/010048
(87) N° publication PCT/PCT Publication No.: 2004/014932
(30) Priorités/Priorities: 2002/08/08 (2002/232074) JP;
2002/11/05 (2002/321729) JP




(51) Cl.Int.⁷/Int.Cl.⁷ C07D 405/12, A61K 31/7056, A61P 9/12,
A61P 9/10, A61P 3/10, A61P 43/00, A61P 19/06,
A61P 3/06, A61P 9/04, A61P 3/04, C07H 17/02

(71) Demandeur/Applicant:
KISSEI PHARMACEUTICAL CO., LTD., JP

(72) Inventeurs/Inventors:
TERANISHI, HIROTAKA, JP;
FUSHIMI, NOBUHIKO, JP;
YONEKUBO, SHIGERU, JP;
SHIMIZU, KAZUO, JP;
SHIBAZAKI, TOSHIHIDE, JP;
ISAJI, MASAYUKI, JP

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : DERIVE DE PYRAZOLE, COMPOSITION MEDICINALE CONTENANT CE DERIVE, UTILISATION
THERAPEUTIQUE DE CEUX-CI ET INTERMEDIAIRE POUR LA PRODUCTION DE CETTE COMPOSITION
(54) Title: PYRAZOLE DERIVATIVE, MEDICINAL COMPOSITION CONTAINING THE SAME, MEDICINAL USE
THEREOF, AND INTERMEDIATE FOR PRODUCTION THEREOF



(57) Abrégé/Abstract:

A pyrazole derivative represented by the general formula (I) (wherein R¹ is H, optionally substituted C₁₋₆ alkyl, etc.; either of Q and T is the group of the formula (II) or the formula (III) and the other is optionally substituted C₁₋₆ alkyl, etc.; R² is H, halogeno, OH, optionally substituted C₁₋₆ alkyl, etc.; X is a single bond, O, or S; Y is a single bond, C₁₋₆ alkylene, etc.; Z is CO or SO₂; R⁴ and R⁵ each is H, optionally substituted C₁₋₆ alkyl, etc.; and R³, R⁶, and R⁷ each is H, halogeno, etc.), a pharmacologically acceptable salt of the derivative, or a prodrug of either. They have excellent human SGLT1 inhibitory activity and are useful as a preventive or therapeutic agent for diseases attributable to hyperglycemia such as diabetes, complications of diabetes, and obesity.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPQ 191


OPIC




CIPO

ABSTRACT


The present invention provides pyrazole derivatives represented by the general formula:



wherein R^1 represents H, an optionally substituted C_{1-6} alkyl group etc.; one of Q and T represents a group represented by the general formula:



10 or a group represented by the general formula:




while the other represents an optionally substituted C_{1-6} alkyl group etc.; R^2 represents H, a halogen atom, OH, an optionally substituted C_{1-6} alkyl group etc.; X represents a single bond, O or S; Y represents a single bond, a C_{1-6} alkylene group etc.; Z represents CO or SO₂; R^4 and R^5 represent H, an optionally substituted C_{1-6} alkyl group etc.; and R^3 , R^6 and R^7 represent

15

H, a halogen atom etc., pharmaceutically acceptable salts thereof, or prodrugs thereof, which exhibit an excellent inhibitory activity in human SGLT1 and are useful as agents for the prevention or treatment of a disease associated with hyperglycemia such as diabetes, diabetic complications or obesity, and
5 pharmaceutical compositions comprising the same, pharmaceutical uses thereof, and intermediates for production thereof.

CLAIMS


1. A pyrazole derivative represented by the general formula:




- 5 wherein

R^1 represents a hydrogen atom, a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a hydroxy(C₂₋₆ alkyl) group, a C₃₋₇ cycloalkyl group, a C₃₋₇ cycloalkyl-substituted (C₁₋₆ alkyl) group, an aryl group which may have the same or different 1 to 3 substituents
 10 selected from the group consisting of a halogen atom, a hydroxy group, an amino group, a C₁₋₆ alkyl group and a C₁₋₆ alkoxy group, or an aryl(C₁₋₆ alkyl) group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group, an amino group, a C₁₋₆ alkyl group and
 15 a C₁₋₆ alkoxy group on the ring;

one of Q and T represents a group represented by the formula:



or a group represented by the formula:



while the other represents a C₁₋₆ alkyl group, a halo(C₁₋₆ alkyl) group, a C₁₋₆ alkoxy-substituted (C₁₋₆ alkyl) group or a C₃₋₇ cycloalkyl group;

5 R² represents a hydrogen atom, a halogen atom, a hydroxy group, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, a C₁₋₆ alkylthio group, a halo(C₁₋₆ alkyl) group, a halo(C₁₋₆ alkoxy) group, a C₁₋₆ alkoxy-substituted (C₁₋₆ alkoxy) group, a C₃₋₇ cycloalkyl-substituted (C₂₋₆ alkoxy) group or a group of the general formula:
 10 -A-R⁸ in which A represents a single bond, an oxygen atom, a methylene group, an ethylene group, -OCH₂- or -CH₂O-; and R⁸ represents a C₃₋₇ cycloalkyl group, a C₂₋₆ heterocycloalkyl group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen
 15 atom, a hydroxy group, an amino group, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, a C₂₋₆ alkenyloxy group, a halo(C₁₋₆ alkyl) group, a hydroxy(C₁₋₆ alkyl) group, a carboxy group, a C₂₋₇ alkoxycarbonyl group, a cyano group and a nitro group, or a heteroaryl group which may have a substituent selected from the
 20 group consisting of a halogen atom and a C₁₋₆ alkyl group;

X represents a single bond, an oxygen atom or a sulfur atom;

Y represents a single bond, a C₁₋₆ alkylene group or a C₂₋₆ alkenylene group with the proviso that X is a single bond
 25 when Y is a single bond;

Z represents a carbonyl group or a sulfonyl group;

R⁴ and R⁵ are the same or different, and each represents a hydrogen atom or a C₁₋₆ alkyl group which may have the same or different 1 to 3 groups selected from the following substituent group (i), or they bind together with the neighboring nitrogen atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group;

R³, R⁶ and R⁷ are the same or different, and each represents a hydrogen atom, a halogen atom, a C₁₋₆ alkyl group or a C₁₋₆ alkoxy group; and

substituent group (i) consists of a hydroxy group, an amino group, a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group, an ureido group, a sulfamide group, a mono or di(C₁₋₆ alkyl)ureido group, a mono or di(C₁₋₆ alkyl)sulfamide group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group, a group of the general formula: -CON(R⁹)R¹⁰ in which R⁹ and R¹⁰ are the same or different, and each represents a hydrogen atom or a C₁₋₆ alkyl group which may have the same or different 1 to 3 substituents selected from the group consisting of a hydroxy group, an amino group, a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group, an ureido group, a mono or di(C₁₋₆ alkyl)ureido group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group and a carbamoyl group, or they bind together with the neighboring nitrogen atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group

consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group,
a C₃₋₇ cycloalkyl group, a C₂₋₆ heterocycloalkyl group, an aryl
group which may have the same or different 1 to 3 substituents
selected from the group consisting of a halogen atom, a hydroxy
5 group, an amino group, a C₁₋₆ alkyl group and a C₁₋₆ alkoxy group,
a heteroaryl group which may have a substituent selected from
the group consisting of a halogen atom and a C₁₋₆ alkyl group,
a C₂₋₆ cyclic amino group which may have a substituent selected
from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆
10 alkyl) group, and a C₁₋₄ aromatic cyclic amino group which may
have a C₁₋₆ alkyl group as a substituent,
or a pharmaceutically acceptable salt thereof.

2. A pyrazole derivative as claimed in claim 1, wherein Y
15 represents a C₁₋₆ alkylene group or a C₂₋₆ alkenylene group;
one of R⁴ and R⁵ represents a C₁₋₆ alkyl group which has the
same or different 1 to 3 groups selected from the following
substituent group (i), the other represents a hydrogen atom or
a C₁₋₆ alkyl group which may have the same or different 1 to
20 3 groups selected from the following substituent group (i); and
substituent group (i) consists of a hydroxy group, an amino group,
a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆
alkyl)]amino group, an ureido group, a sulfamide group, a mono
or di(C₁₋₆ alkyl)ureido group, a mono or di(C₁₋₆ alkyl)sulfamide
25 group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group,
a group of the general formula: -CON(R⁹)R¹⁰ in which R⁹ and R¹⁰
are the same or different, and each represents a hydrogen atom

or a C₁₋₆ alkyl group which may have the same or different 1 to 3 substituents selected from the group consisting of a hydroxy group, an amino group, a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group, an ureido group, 5 a mono or di(C₁₋₆ alkyl)ureido group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group and a carbamoyl group, or they bind together with the neighboring nitrogen atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ 10 alkyl) group, a C₃₋₇ cycloalkyl group, a C₂₋₆ heterocycloalkyl group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group, an amino group, a C₁₋₆ alkyl group and a C₁₋₆ alkoxy group, a heteroaryl group which may have a 15 substituent selected from the group consisting of a halogen atom and a C₁₋₆ alkyl group, a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group, and a C₁₋₄ aromatic cyclic amino group which may have a C₁₋₆ alkyl group as a substituent, 20 or a pharmaceutically acceptable salt thereof.

3. A pyrazole derivative as claimed in claim 2, wherein one of R⁴ and R⁵ represents a C₁₋₆ alkyl group which has a group selected from the following substituent group (iA), the other 25 represents a hydrogen atom; and substituent group (iA) is a group of the general formula: -CON(R^{9A})R^{10A} in which R^{9A} and R^{10A} bind together with the neighboring nitrogen atom to form a C₂₋₆ cyclic

amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group, or a pharmaceutically acceptable salt thereof.

5 4. A pyrazole derivative as claimed in any one of claims 1-3, wherein X represents a single bond; and Y represents a trimethylene group or a 1-propenylene group, or a pharmaceutically acceptable salt thereof.


10 5. A pyrazole derivative as claimed in any one of claims 1-3, wherein X represents an oxygen atom; and Y represents an ethylene group or a trimethylene group, or a pharmaceutically acceptable salt thereof.

15 6. A pyrazole derivative as claimed in claim 1, wherein X represents a single bond; Y represents a single bond; one of R⁴ and R⁵ represents a C₁₋₆ alkyl group which has the same or different 1 to 3 groups selected from the following substituent group (iB), the other represents a hydrogen atom or a C₁₋₆ alkyl
20 group which may have the same or different 1 to 3 groups selected from the following substituent group (iB); and substituent group (iB) consists of an ureido group, a sulfamide group, a mono or di(C₁₋₆ alkyl)ureido group, a mono or di(C₁₋₆ alkyl)sulfamide group, a C₁₋₆ alkylsulfonylamino group, a group of the general
25 formula: -CON(R^{9B})R^{10B} in which one of R^{9B} and R^{10B} represents a C₁₋₆ alkyl group which has the same or different 1 to 3 substituents selected from the group consisting of a hydroxy


group, an amino group, a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group, an ureido group, a mono or di(C₁₋₆ alkyl)ureido group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group and a carbamoyl group, the other
5 represents a hydrogen atom, a C₁₋₆ alkyl group which may have the same or different 1 to 3 substituents selected from the group consisting of a hydroxy group, an amino group, a mono or di(C₁₋₆ alkyl)amino group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group, an ureido group, a mono or di(C₁₋₆ alkyl)ureido group, a C₂₋₇
10 acylamino group, a C₁₋₆ alkylsulfonylamino group and a carbamoyl group, or they bind together with the neighboring nitrogen atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group, a C₃₋₇ cycloalkyl group, a C₂₋₆
15 heterocycloalkyl group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group, an amino group, a C₁₋₆ alkyl group and a C₁₋₆ alkoxy group, a heteroaryl group which may have a substituent selected from the group consisting
20 of a halogen atom and a C₁₋₆ alkyl group, a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group, and a C₁₋₄ aromatic cyclic amino group which may have a C₁₋₆ alkyl group as a substituent, or a pharmaceutically acceptable
25 salt thereof.

7. A pyrazole derivative as claimed in any one of claims 1-6,

wherein R^1 represents a hydrogen atom or a hydroxy(C_{2-6} alkyl) group; T represents a group represented by the formula:




or a group represented by the formula:



5


Q represents a C_{1-6} alkyl group or a halo(C_{1-6} alkyl) group; and R^3 , R^6 and R^7 represent a hydrogen atom, or a pharmaceutically acceptable salt thereof.

- 10 8. A pyrazole derivative as claimed in any one of claims 1-6, wherein one of Q and T represents a group represented by the formula:



- the other represents a C_{1-6} alkyl group, a halo(C_{1-6} alkyl) group,
15 a C_{1-6} alkoxy-substituted (C_{1-6} alkyl) group or a C_{3-7} cycloalkyl group, or a pharmaceutically acceptable salt thereof.

9. A pyrazole derivative as claimed in claim 7 or 8, wherein T represents a group represented by the formula:




or a pharmaceutically acceptable salt thereof.

10. A pyrazole derivative as claimed in claim 7 or 9, wherein
 5 Q represents an isopropyl group, or a pharmaceutically acceptable salt thereof.


11. A prodrug of a pyrazole derivative as claimed in any one of claims 1-10 or a pharmaceutically acceptable salt thereof.

10

12. A prodrug as claimed in claim 11, wherein T represents a group represented by the formula:



or a group represented by the formula:



15

in which the hydroxy group at the 4-position is substituted by a glucopyranosyl group or a galactopyranosyl group, or the hydroxy group at the 6-position is substituted by a glucopyranosyl group, a galactopyranosyl group, a C₂₋₇ acyl group,

a C₁₋₆ alkoxy-substituted (C₂₋₇ acyl) group, a C₂₋₇ alkoxy-carbonyl-substituted (C₂₋₇ acyl) group, a C₂₋₇ alkoxycarbonyl group, an aryl(C₂₋₇ alkoxycarbonyl) group or a C₁₋₆ alkoxy-substituted (C₂₋₇ alkoxycarbonyl) group.

5

13. A pyrazole derivative as claimed in claim 1, which is a compound selected from the following group:

4-[(4-{3-[1-carbamoyl-1-(methyl)ethylcarbamoyl]propyl}-2-methylphenyl)methyl]-3-(β-D-glucopyranosyloxy)-5-isopropyl-
10 1H-pyrazole;

3-(β-D-galactopyranosyloxy)-4-[(4-{3-[1-{[4-(2-hydroxyethyl)piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]-propyl}phenyl)methyl]-5-isopropyl-1H-pyrazole;

3-(β-D-galactopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-{2-(dimethylamino)ethylcarbamoyl}-1-(methyl)ethylcarbamoyl]-propyl}phenyl)methyl]-1H-pyrazole;

4-[(4-{3-[1-(2-aminoethylcarbamoyl)-1-(methyl)ethylcarbamoyl]propyl}phenyl)methyl]-3-(β-D-galactopyranosyloxy)-5-isopropyl-1H-pyrazole;

20 3-(β-D-galactopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-propyl}phenyl)methyl]-1H-pyrazole;

3-(β-D-glucopyranosyloxy)-4-[(4-{3-[1-{[4-(2-hydroxyethyl)piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]-propyl}-2-methylphenyl)methyl]-5-isopropyl-1H-pyrazole;

25 3-(β-D-galactopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-[(4-methylpiperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-

propyl)phenyl)methyl}-1H-pyrazole;

3-(β -D-galactopyranosyloxy)-5-isopropyl-4-([4-(3-{1-[(4-isopropylpiperazin-1-yl)carbonyl]-1-(methyl)ethyl-carbamoyl}propyl)phenyl)methyl}-1H-pyrazole;

5 3-(β -D-glucopyranosyloxy)-4-[(4-{3-[(S)-2-hydroxy-1-(methyl)ethylcarbamoyl]propyl)phenyl)methyl]-5-isopropyl-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-4-[(4-{(1E)-3-[(S)-2-hydroxy-1-(methyl)ethylcarbamoyl]prop-1-enyl}phenyl)methyl]-5-

10 isopropyl-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-5-isopropyl-4-([4-(2-{1-[(4-methylpiperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-ethoxy}-2-methylphenyl)methyl)-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-4-[(4-{2-[2-hydroxy-1,1-di-(methyl)ethylcarbamoyl]ethoxy}-2-methylphenyl)methyl]-5-
15 isopropyl-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-4-[(4-{2-[1-[(4-(2-hydroxyethyl)-piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]ethoxy}-2-methylphenyl)methyl]-5-isopropyl-1H-pyrazole;

20 3-(β -D-glucopyranosyloxy)-5-isopropyl-4-([4-(2-{1-[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-ethoxy}-2-methylphenyl)methyl)-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-5-isopropyl-4-([4-(3-{1-[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-propyl)-2-methylphenyl)methyl)-1H-pyrazole;

25 3-(β -D-glucopyranosyloxy)-5-isopropyl-4-([4-(3-{1-[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-

- propoxy)-2-methylphenyl)methyl]-1H-pyrazole;
 3-(β -D-glucopyranosyloxy)-4-[(4-{3-[1-{[4-(2-hydroxyethyl)-
 piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]propoxy)-
 2-methylphenyl)methyl]-5-isopropyl-1H-pyrazole;
 5 3-(β -D-glucopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-
 [(4-methylpiperazin-1-yl)carbonyl]-1-(methyl)ethyl-
 carbamoyl]propoxy)-2-methylphenyl)methyl]-1H-pyrazole;
 3-(β -D-galactopyranosyloxy)-1-(3-hydroxypropyl)-5-
 isopropyl-4-[(4-{3-[1-[(piperazin-1-yl)carbonyl]-1-
 10 (methyl)ethylcarbamoyl]propyl)phenyl)methyl]-1H-pyrazole;
 3-(β -D-galactopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-
 [(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
 propoxy)-2-methylphenyl)methyl]-1H-pyrazole;
 4-[(2-fluoro-4-{3-[1-[(piperazin-1-yl)carbonyl]-1-(methyl)-
 15 ethylcarbamoyl]propyl)phenyl)methyl]-3-(β -D-galacto-
 pyranosyloxy)-5-isopropyl-1H-pyrazole;
 4-[(2-chloro-4-{3-[1-[(piperazin-1-yl)carbonyl]-1-(methyl)-
 ethylcarbamoyl]propyl)phenyl)methyl]-3-(β -D-glucopyranosyl-
 oxy)-5-isopropyl-1H-pyrazole, and
 20 pharmaceutically acceptable salts thereof.

14. A pyrazole derivative as claimed in claim 13, which is
 a compound selected from the following group:

- 3-(β -D-galactopyranosyloxy)-4-[(4-{3-[1-{[4-(2-hydroxy-
 25 ethyl)piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]-
 propyl)phenyl)methyl]-5-isopropyl-1H-pyrazole;
 3-(β -D-galactopyranosyloxy)-5-isopropyl-4-[(4-{3-[1-

[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl}-
propyl)phenyl)methyl]-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-4-[(4-{3-[1-{[4-(2-hydroxyethyl)-
piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]-

5 propyl)-2-methylphenyl)methyl]-5-isopropyl-1H-pyrazole;

3-(β -D-galactopyranosyloxy)-5-isopropyl-4-[[4-(3-{1-[(4-
methylpiperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
propyl)phenyl)methyl]-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-5-isopropyl-4-[[4-(2-{1-[(4-
10 methylpiperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
ethoxy)-2-methylphenyl)methyl]-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-4-[(4-{2-[1-{[4-(2-hydroxyethyl)-
piperazin-1-yl]carbonyl}-1-(methyl)ethylcarbamoyl]ethoxy}-
2-methylphenyl)methyl]-5-isopropyl-1H-pyrazole;

15 3-(β -D-glucopyranosyloxy)-5-isopropyl-4-[[4-(2-{1-
[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
ethoxy)-2-methylphenyl)methyl]-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-5-isopropyl-4-[[4-(3-{1-
[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-

20 propyl)-2-methylphenyl)methyl]-1H-pyrazole;

3-(β -D-glucopyranosyloxy)-5-isopropyl-4-[[4-(3-{1-
[(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
propoxy)-2-methylphenyl)methyl]-1H-pyrazole;

3-(β -D-galactopyranosyloxy)-5-isopropyl-4-[[4-(3-{1-
25 [(piperazin-1-yl)carbonyl]-1-(methyl)ethylcarbamoyl]-
propoxy)-2-methylphenyl)methyl]-1H-pyrazole;

4-[[2-fluoro-4-(3-{1-[(piperazin-1-yl)carbonyl]-1-(methyl)-

ethylcarbamoyl}propyl}phenyl}methyl}-3-(β -D-galactopyranosyloxy)-5-isopropyl-1H-pyrazole, and pharmaceutically acceptable salts thereof.

- 5 15. A pharmaceutical composition comprising as an active ingredient a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.
- 10 16. A human SGLT1 inhibitor comprising as an active ingredient a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.
- 15 17. An agent for inhibiting postprandial hyperglycemia comprising as an active ingredient a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.
- 20 18. An agent for the prevention or treatment of a disease associated with hyperglycemia, which comprises as an active ingredient a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.
- 25 19. An agent for the prevention or treatment as claimed in claim 18, wherein the disease associated with hyperglycemia is a disease selected from the group consisting of diabetes,

impaired glucose tolerance, diabetic complications, obesity,
hyperinsulinemia, hyperlipidemia, hypercholesterolemia,
hypertriglyceridemia, lipid metabolism disorder,
atherosclerosis, hypertension, congestive heart failure, edema,
5 hyperuricemia and gout.

20. An agent for the inhibition of advancing impaired glucose
tolerance or impaired fasting glycemia into diabetes in a subject,
which comprises as an active ingredient a pyrazole derivative
10 as claimed in any one of claims 1-14, a pharmaceutically
acceptable salt thereof or a prodrug thereof.

21. An agent for the prevention or treatment of a disease
associated with the increase of blood galactose level, which
15 comprises as an active ingredient a pyrazole derivative as
claimed in any one of claims 1-14, a pharmaceutically acceptable
salt thereof or a prodrug thereof.

22. An agent for the prevention or treatment as claimed in
20 claim 21, wherein the disease associated with the increase of
blood galactose level is galactosemia.

23. A pharmaceutical composition as claimed in claim 15,
wherein the dosage form is sustained release formulation.

25

24. An agent as claimed in any one of claims 16-22, wherein
the dosage form is sustained release formulation.

25. A method for the prevention or treatment of a disease associated with hyperglycemia, which comprises administering an effective amount of a pyrazole derivative as claimed in any
5 one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.

26. A method for the inhibition of advancing impaired glucose tolerance into diabetes in a subject, which comprises
10 administering an effective amount of a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof.

27. A use of a pyrazole derivative as claimed in any one of
15 claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof for the manufacture of a pharmaceutical composition for the prevention or treatment of a disease associated with hyperglycemia.

20 28. A use of a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof for the manufacture of a pharmaceutical composition for the inhibition of advancing impaired glucose tolerance into diabetes in a subject.

25

29. A pharmaceutical combination which comprises (A) a pyrazole derivative as claimed in any one of claims 1-14, a

pharmaceutically acceptable salt thereof or a prodrug thereof,
and (B) at least one member selected from the group consisting
of an insulin sensitivity enhancer, a glucose absorption
inhibitor, a biguanide, an insulin secretion enhancer, a SGLT2
5 inhibitor, an insulin or insulin analogue, a glucagon receptor
antagonist, an insulin receptor kinase stimulant, a tripeptidyl
peptidase II inhibitor, a dipeptidyl peptidase IV inhibitor,
a protein tyrosine phosphatase-1B inhibitor, a glycogen
phosphorylase inhibitor, a glucose-6-phosphatase inhibitor, a
10 fructose-bisphosphatase inhibitor, a pyruvate dehydrogenase
inhibitor, a hepatic gluconeogenesis inhibitor, D-chiroinsitol,
a glycogen synthase kinase-3 inhibitor, glucagon-like peptide-1,
a glucagon-like peptide-1 analogue, a glucagon-like peptide-1
agonist, amylin, an amylin analogue, an amylin agonist, an aldose
15 reductase inhibitor, an advanced glycation endproducts
formation inhibitor, a protein kinase C inhibitor, a
 γ -aminobutyric acid receptor antagonist, a sodium channel
antagonist, a transcript factor NF- κ B inhibitor, a lipid
peroxidase inhibitor, an *N*-acetylated- α -linked-acid-
20 dipeptidase inhibitor, insulin-like growth factor-I,
platelet-derived growth factor, a platelet-derived growth
factor analogue, epidermal growth factor, nerve growth factor,
a carnitine derivative, uridine, 5-hydroxy-1-methylhidantoin,
EGB-761, bimoclomol, sulodexide, Y-128, antidiarrhoics,
25 cathartics, a hydroxymethylglutaryl coenzyme A reductase
inhibitor, a fibric acid derivative, a β_3 -adrenoceptor agonist,
an acyl-coenzyme A cholesterol acyltransferase inhibitor,

probcol, a thyroid hormone receptor agonist, a cholesterol absorption inhibitor, a lipase inhibitor, a microsomal triglyceride transfer protein inhibitor, a lipoxygenase inhibitor, a carnitine palmitoyl-transferase inhibitor, a squalene synthase inhibitor, a low-density lipoprotein receptor enhancer, a nicotinic acid derivative, a bile acid sequestrant, a sodium/bile acid cotransporter inhibitor, a cholesterol ester transfer protein inhibitor, an appetite suppressant, an angiotensin-converting enzyme inhibitor, a neutral endopeptidase inhibitor, an angiotensin II receptor antagonist, an endothelin-converting enzyme inhibitor, an endothelin receptor antagonist, a diuretic agent, a calcium antagonist, a vasodilating antihypertensive agent, a sympathetic blocking agent, a centrally acting antihypertensive agent, an α_2 -adrenoceptor agonist, an antiplatelets agent, a uric acid synthesis inhibitor, a uricosuric agent and a urinary alkalinizer.

30. A method for the prevention or treatment of a disease associated with hyperglycemia, which comprises administering an effective amount of (A) a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof, and (B) at least one member selected from the group consisting of an insulin sensitivity enhancer, a glucose absorption inhibitor, a biguanide, an insulin secretion enhancer, a SGLT2 inhibitor, an insulin or insulin analogue, a glucagon receptor antagonist, an insulin receptor kinase

stimulant, a tripeptidyl peptidase II inhibitor, a dipeptidyl peptidase IV inhibitor, a protein tyrosine phosphatase-1B inhibitor, a glycogen phosphorylase inhibitor, a glucose-6-phosphatase inhibitor, a fructose-bisphosphatase inhibitor, a pyruvate dehydrogenase inhibitor, a hepatic gluconeogenesis inhibitor, D-chiroinsitol, a glycogen synthase kinase-3 inhibitor, glucagon-like peptide-1, a glucagon-like peptide-1 analogue, a glucagon-like peptide-1 agonist, amylin, an amylin analogue, an amylin agonist, an aldose reductase inhibitor, an advanced glycation endproducts formation inhibitor, a protein kinase C inhibitor, a γ -aminobutyric acid receptor antagonist, a sodium channel antagonist, a transcript factor NF- κ B inhibitor, a lipid peroxidase inhibitor, an N-acetylated- α -linked-acid-dipeptidase inhibitor, insulin-like growth factor-I, platelet-derived growth factor, a platelet-derived growth factor analogue, epidermal growth factor, nerve growth factor, a carnitine derivative, uridine, 5-hydroxy-1-methylhydantoin, EGB-761, bimoclomol, sulodexide, Y-128, antidiarrhoics, cathartics, a hydroxymethylglutaryl coenzyme A reductase inhibitor, a fibric acid derivative, a β_3 -adrenoceptor agonist, an acyl-coenzyme A cholesterol acyltransferase inhibitor, probcol, a thyroid hormone receptor agonist, a cholesterol absorption inhibitor, a lipase inhibitor, a microsomal triglyceride transfer protein inhibitor, a lipoxxygenase inhibitor, a carnitine palmitoyl-transferase inhibitor, a squalene synthase inhibitor, a low-density lipoprotein receptor enhancer, a nicotinic acid derivative, a

bile acid sequestrant, a sodium/bile acid cotransporter inhibitor, a cholesterol ester transfer protein inhibitor, an appetite suppressant, an angiotensin-converting enzyme inhibitor, a neutral endopeptidase inhibitor, an angiotensin
5 II receptor antagonist, an endothelin-converting enzyme inhibitor, an endothelin receptor antagonist, a diuretic agent, a calcium antagonist, a vasodilating antihypertensive agent, a sympathetic blocking agent, a centrally acting antihypertensive agent, an α_2 -adrenoceptor agonist, an
10 antiplatelets agent, a uric acid synthesis inhibitor, a uricosuric agent and a urinary alkalinizer.

31. A method for the inhibition of advancing impaired glucose tolerance into diabetes in a subject, which comprises
15 administering an effective amount of (A) a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof, and (B) at least one member selected from the group consisting of an insulin sensitivity enhancer, a glucose absorption inhibitor, a
20 biguanide, an insulin secretion enhancer, a SGLT2 inhibitor, an insulin or insulin analogue, a glucagon receptor antagonist, an insulin receptor kinase stimulant, a tripeptidyl peptidase II inhibitor, a dipeptidyl peptidase IV inhibitor, a protein tyrosine phosphatase-1B inhibitor, a glycogen phosphorylase
25 inhibitor, a glucose-6-phosphatase inhibitor, a fructose-bisphosphatase inhibitor, a pyruvate dehydrogenase inhibitor, a hepatic gluconeogenesis inhibitor, D-chiroinsitol,

a glycogen synthase kinase-3 inhibitor, glucagon-like peptide-1, a glucagon-like peptide-1 analogue, a glucagon-like peptide-1 agonist, amylin, an amylin analogue, an amylin agonist, an aldose reductase inhibitor, an advanced glycation endproducts
5 formation inhibitor, a protein kinase C inhibitor, a γ -aminobutyric acid receptor antagonist, a sodium channel antagonist, a transcript factor NF- κ B inhibitor, a lipid peroxidase inhibitor, an N-acetylated- α -linked-acid-dipeptidase inhibitor, insulin-like growth factor-I,
10 platelet-derived growth factor, a platelet-derived growth factor analogue, epidermal growth factor, nerve growth factor, a carnitine derivative, uridine, 5-hydroxy-1-methylhydantoin, EGB-761, bimoclomol, sulodexide, Y-128, antidiarrhoics, cathartics, a hydroxymethylglutaryl coenzyme A reductase
15 inhibitor, a fibrin acid derivative, a β_3 -adrenoceptor agonist, an acyl-coenzyme A cholesterol acyltransferase inhibitor, probucol, a thyroid hormone receptor agonist, a cholesterol absorption inhibitor, a lipase inhibitor, a microsomal triglyceride transfer protein inhibitor, a lipoxygenase
20 inhibitor, a carnitine palmitoyl-transferase inhibitor, a squalene synthase inhibitor, a low-density lipoprotein receptor enhancer, a nicotinic acid derivative, a bile acid sequestrant, a sodium/bile acid cotransporter inhibitor, a cholesterol ester transfer protein inhibitor, an appetite suppressant, an
25 angiotensin-converting enzyme inhibitor, a neutral endopeptidase inhibitor, an angiotensin II receptor antagonist, an endothelin-converting enzyme inhibitor, an endothelin

receptor antagonist, a diuretic agent, a calcium antagonist, a vasodilating antihypertensive agent, a sympathetic blocking agent, a centrally acting antihypertensive agent, an α_2 -adrenoceptor agonist, an antiplatelets agent, a uric acid synthesis inhibitor, a uricosuric agent and a urinary alkalinizer.

32. A use of (A) a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof, and (B) at least one member selected from the group consisting of an insulin sensitivity enhancer, a glucose absorption inhibitor, a biguanide, an insulin secretion enhancer, a SGLT2 inhibitor, an insulin or insulin analogue, a glucagon receptor antagonist, an insulin receptor kinase stimulant, a tripeptidyl peptidase II inhibitor, a dipeptidyl peptidase IV inhibitor, a protein tyrosine phosphatase-1B inhibitor, a glycogen phosphorylase inhibitor, a glucose-6-phosphatase inhibitor, a fructose-bisphosphatase inhibitor, a pyruvate dehydrogenase inhibitor, a hepatic gluconeogenesis inhibitor, D-chiroinsitol, a glycogen synthase kinase-3 inhibitor, glucagon-like peptide-1, a glucagon-like peptide-1 analogue, a glucagon-like peptide-1 agonist, amylin, an amylin analogue, an amylin agonist, an aldose reductase inhibitor, an advanced glycation endproducts formation inhibitor, a protein kinase C inhibitor, a γ -aminobutyric acid receptor antagonist, a sodium channel antagonist, a transcript factor NF- κ B inhibitor, a lipid peroxidase inhibitor, an N-acetylated- α -linked-acid-

dipeptidase inhibitor, insulin-like growth factor-I, platelet-derived growth factor, a platelet-derived growth factor analogue, epidermal growth factor, nerve growth factor, a carnitine derivative, uridine, 5-hydroxy-1-methylhydantoin, 5 EGB-761, bimoclomol, sulodexide, Y-128, antidiarrhoics, cathartics, a hydroxymethylglutaryl coenzyme A reductase inhibitor, a fibric acid derivative, a β_3 -adrenoceptor agonist, an acyl-coenzyme A cholesterol acyltransferase inhibitor, probcol, a thyroid hormone receptor agonist, a cholesterol 10 absorption inhibitor, a lipase inhibitor, a microsomal triglyceride transfer protein inhibitor, a lipoxygenase inhibitor, a carnitine palmitoyl-transferase inhibitor, a squalene synthase inhibitor, a low-density lipoprotein receptor enhancer, a nicotinic acid derivative, a bile acid sequestrant, 15 a sodium/bile acid cotransporter inhibitor, a cholesterol ester transfer protein inhibitor, an appetite suppressant, an angiotensin-converting enzyme inhibitor, a neutral endopeptidase inhibitor, an angiotensin II receptor antagonist, an endothelin-converting enzyme inhibitor, an endothelin 20 receptor antagonist, a diuretic agent, a calcium antagonist, a vasodilating antihypertensive agent, a sympathetic blocking agent, a centrally acting antihypertensive agent, an α_2 -adrenoceptor agonist, an antiplatelets agent, a uric acid synthesis inhibitor, auricosuric agent and a urinary alkalinizer, 25 for the manufacture of a pharmaceutical composition for the prevention or treatment of a disease associated with hyperglycemia.

33. A use of (A) a pyrazole derivative as claimed in any one of claims 1-14, a pharmaceutically acceptable salt thereof or a prodrug thereof, and (B) at least one member selected from
5 the group consisting of an insulin sensitivity enhancer, a glucose absorption inhibitor, a biguanide, an insulin secretion enhancer, a SGLT2 inhibitor, an insulin or insulin analogue, a glucagon receptor antagonist, an insulin receptor kinase stimulant, a tripeptidyl peptidase II inhibitor, a dipeptidyl
10 peptidase IV inhibitor, a protein tyrosine phosphatase-1B inhibitor, a glycogen phosphorylase inhibitor, a glucose-6-phosphatase inhibitor, a fructose-bisphosphatase inhibitor, a pyruvate dehydrogenase inhibitor, a hepatic gluconeogenesis inhibitor, D-chiroinsitol, a glycogen synthase kinase-3
15 inhibitor, glucagon-like peptide-1, a glucagon-like peptide-1 analogue, a glucagon-like peptide-1 agonist, amylin, an amylin analogue, an amylin agonist, an aldose reductase inhibitor, an advanced glycation endproducts formation inhibitor, a protein kinase C inhibitor, a γ -aminobutyric acid receptor antagonist,
20 asodiumchannel antagonist, a transcript factor NF- κ B inhibitor, a lipid peroxidase inhibitor, an *N*-acetylated- α -linked-acid-dipeptidase inhibitor, insulin-like growth factor-I, platelet-derived growth factor, a platelet-derived growth factor analogue, epidermal growth factor, nerve growth factor,
25 a carnitine derivative, uridine, 5-hydroxy-1-methylhidantoin, EGB-761, bimoclomol, sulodexide, Y-128, antidiarrhoics, cathartics, a hydroxymethylglutaryl coenzyme A reductase

inhibitor, a fibric acid derivative, a β_3 -adrenoceptor agonist, an acyl-coenzyme A cholesterol acyltransferase inhibitor, probcol, a thyroid hormone receptor agonist, a cholesterol absorption inhibitor, a lipase inhibitor, a microsomal


5 triglyceride transfer protein inhibitor, a lipoxxygenase inhibitor, a carnitine palmitoyl-transferase inhibitor, a squalene synthase inhibitor, a low-density lipoprotein receptor enhancer, a nicotinic acid derivative, a bile acid sequestrant, a sodium/bile acid cotransporter inhibitor, a cholesterol ester

10 transfer protein inhibitor, an appetite suppressant, an angiotensin-converting enzyme inhibitor, a neutral endopeptidase inhibitor, an angiotensin II receptor antagonist, an endothelin-converting enzyme inhibitor, an endothelin receptor antagonist, a diuretic agent, a calcium antagonist,

15 a vasodilating antihypertensive agent, a sympathetic blocking agent, a centrally acting antihypertensive agent, an α_2 -adrenoceptor agonist, an antiplatelets agent, a uric acid synthesis inhibitor, a uricosuric agent and a urinary alkalinizer, for the manufacture of a pharmaceutical composition for the

20 inhibition of advancing impaired glucose tolerance into diabetes in a subject.

34. A pyrazole derivative represented by the general formula:



wherein

R^{11} represents a hydrogen atom, a C_{1-6} alkyl group, a C_{2-6} alkenyl group, a hydroxy(C_{2-6} alkyl) group which may have a
 5 protective group, a C_{3-7} cycloalkyl group, a C_{3-7} cycloalkyl-substituted (C_{1-6} alkyl) group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group which may have a protective group, an amino group which may have a protective
 10 group, a C_{1-6} alkyl group and a C_{1-6} alkoxy group, or an aryl(C_{1-6} alkyl) group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group which may have a protective group, an amino group which may have a protective group, a C_{1-6} alkyl group and
 15 a C_{1-6} alkoxy group on the ring;

one of Q^2 and T^2 represents a 2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyloxy group or a 2,3,4,6-tetra-*O*-acetyl- β -D-galactopyranosyloxy group, while the other represents a C_{1-6} alkyl group, a halo(C_{1-6} alkyl) group, a C_{1-6} alkoxy-substituted
 20 (C_{1-6} alkyl) group or a C_{3-7} cycloalkyl group;

R^{12} represents a hydrogen atom, a halogen atom, a hydroxy group which may have a protective group, a C_{1-6} alkyl group,

a C₁₋₆ alkoxy group, a C₁₋₆ alkylthio group, a halo(C₁₋₆ alkyl) group, a halo(C₁₋₆ alkoxy) group, a C₁₋₆ alkoxy-substituted (C₁₋₆ alkoxy) group, a C₃₋₇ cycloalkyl-substituted (C₂₋₆ alkoxy) group or a group of the general formula: -A-R¹⁸ in which A represents
5 a single bond, an oxygen atom, a methylene group, an ethylene group, -OCH₂- or -CH₂O-; and R¹⁸ represents a C₃₋₇ cycloalkyl group, a C₂₋₆ heterocycloalkyl group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group which
10 may have a protective group, an amino group which may have a protective group, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, a C₂₋₆ alkenyloxy group, a halo(C₁₋₆ alkyl) group, a hydroxy(C₁₋₆ alkyl) group which may have a protective group, a carboxy group which may have a protective group, a C₂₋₇ alkoxycarbonyl group,
15 a cyano group and a nitro group, or a heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and a C₁₋₆ alkyl group;

X represents a single bond, an oxygen atom or a sulfur atom;

20 Y represents a single bond, a C₁₋₆ alkylene group or a C₂₋₆ alkenylene group with the proviso that X is a single bond when Y is a single bond;

Z represents a carbonyl group or a sulfonyl group;

R¹⁴ and R¹⁵ are the same or different, and each represents
25 a hydrogen atom or a C₁₋₆ alkyl group which may have the same or different 1 to 3 groups selected from the following substituent group (ii), or they bind together with the neighboring nitrogen

atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group which may have a protective group;

R³, R⁶ and R⁷ are the same or different, and each represents
5 a hydrogen atom, a halogen atom, a C₁₋₆ alkyl group or a C₁₋₆ alkoxy group; and

substituent group (ii) consists of a hydroxy group which may have a protective group, an amino group which may have a protective group, a mono or di(C₁₋₆ alkyl)amino group which may
10 have a protective group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group which may have a protective group, an ureido group, a sulfamide group, a mono or di(C₁₋₆ alkyl)ureido group, a mono or di(C₁₋₆ alkyl)sulfamide group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group, a group of the general formula:
15 -CON(R¹⁹)R²⁰ in which R¹⁹ and R²⁰ are the same or different, and each represents a hydrogen atom or a C₁₋₆ alkyl group which may have the same or different 1 to 3 substituents selected from the group consisting of a hydroxy group which may have a protective group, an amino group which may have a protective group, a mono
20 or di(C₁₋₆ alkyl)amino group which may have a protective group, a mono or di[hydroxy(C₁₋₆ alkyl)]amino group which may have a protective group, an ureido group, a mono or di(C₁₋₆ alkyl)ureido group, a C₂₋₇ acylamino group, a C₁₋₆ alkylsulfonylamino group and a carbamoyl group, or they bind together with the neighboring
25 nitrogen atom to form a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl group and a hydroxy(C₁₋₆ alkyl) group which may have a protective

group, a C₃₋₇ cycloalkyl group, a C₂₋₆ heterocycloalkyl group, an aryl group which may have the same or different 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxy group which may have a protective group, an amino
5 group which may have a protective group, a C₁₋₆ alkyl group and a C₁₋₆ alkoxy group, a heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and a C₁₋₆ alkyl group, a C₂₋₆ cyclic amino group which may have a substituent selected from the group consisting of a C₁₋₆ alkyl
10 group and a hydroxy(C₁₋₆ alkyl) group which may have a protective group, and a C₁₋₄ aromatic cyclic amino group which may have a C₁₋₆ alkyl group as a substituent, or a salt thereof.

